GOLDSUITE

MAXIMA

goldsuite

KELP IS PACKED WITH OVER 70 VITAMINS, TRACE MINERALS, ENZYMES AND GROWTH HORMONES. MANY OF THESE TRACE MINERALS ARE NOT FOUND ANYWHERE ELSE BUT THE OCEAN. KELP IS 100% NATURAL AND CONTAINS NO HARMFUL CHEMICALS.

BENEFITS OF GOLDSUITE MAXIMA:

- Promotes more efficient uptake of available nutrients from the soil, increasing protein content.
- Increases the number of chloroplasts in the plant which means more green pigment chlorophyll is available for photosynthesis.
- Contains carbohydrates which means stronger roots and bigger leaves.
- Increases resilience to plant stress events, such as frost, fungus, disease and transplantation.
- Roots grow faster and develop more mass.
- Promotes root growth and invigorates early seedling development.

MIXING

- Shake the container thoroughly before use.
- Follow the standard tank mixing procedure if this product is applied in combination with other agricultural products.
- Ensure good agitation throughout application.

COMPATIBILITY

- Goldsuite Maxima is compatible with most agricultural remedies.
- It is however advisable to do a miscibility test prior to mixing Goldsuite Maxima with other agricultural products.

DIRECTIONS FOR USE (USE ONLY AS DIRECTED). Use lower rates as maintenance and higher rates to correct deficiencies.

CROP		RATE / ha	COMMENTS
Berries	i. Blueberries, Blackberries, Raspberries	1 - 3 L 1 - 3 L	i. Apply at the start of the growing season ii. Repeat 2 – 3 times with 14 day intervals Apply in 300 L water/ha
	ii. Strawberries	1 - 2 L 1 - 2 L	i. Drench seedlings before transplanting (see drench application below) ii. Apply 2 weeks after transplant iii. Repeat 14 days later, repeat 2 – 3 times Apply in 200 L water
Bulb, Tuber & Root Crops			
	i. Onions, Garlic	1 - 2 L 1 - 2,5 L	i. Drench seedlings before transplanting (see drench application below) ii. Apply 2 weeks after transplant iii. Repeat 14 days later, repeat 2 – 3 times Apply in 300 - 500 L water
	ii. Potatoes, Sweet Potatoes	1 - 2 L 1 - 2 L 3 - 5 L 3 - 5 L	i. Start of emergence ii. Apply 3 weeks after transplant iii. Repeat 14 days after transplant iv. Repeat 14 days later, repeat 2 – 3 times Apply in 200 L water
	iii. Beetroot, Carrots	1 - 2L 1 - 2,5 L	i. Drench seedlings before transplanting (see drench application below) ii. Apply 2 weeks after transplant iii. Repeat 14 days later, repeat 2 – 3times Apply in 200 L water
Curcubits	Cucumbers, Marrows, Melons, Pumpkins, Baby marrows	1 - 2 L 1 - 2,5 L	i. Drench seedlings before transplanting (see drench application below) ii. Apply 2 weeks after transplant iii. Repeat 14 days later, repeat 2 – 3times Apply in 200 L water
Grains	i. Maize, Sorghum	1 - 2 L 1 - 2,5 L	i. Apply 2 – 3 weeks after emergence ii. Apply 14 days later Apply in 200 L water/ha
	ii. Barley, Oats, Wheat	1 - 3 L 1 - 3 L	i. Apply 2 – 3 weeks after emergence ii. Apply 14 days later Apply in 200 L water/ha
Grapes	Table and Wine Grapes	1 - 3 L 1 - 3 L 1 - 3 L	i. Apply just before flowering ii. Apply at 100% petal drop iii. Apply at pea berry size Apply in 500 - 750 L water/ha
Leafy Vegetables	Lettuce, Spinach	1 - 2L 1 - 2,5 L	i. Drench seedlings before transplanting (see drench application below) ii. Apply 2 weeks after transplant iii. Repeat 14 days later, repeat 2 – 3times Apply in 200 L water
Legumes	i. Dry Beans, Peas	1 - 3 L 1 - 3 L	i. Apply 2 – 3 weeks after emergence ii. Apply 14 days later Apply in 200 L water/ha
Nut Trees	Macadamia, Pecan, Wall nuts	3 - 5 L 3 - 5 L	i. Apply 1 - 2 weeks after bud break ii. Repeat 14 days later iii. Repeat 3 times with 30 day intervals Apply in 1000 - 2000 L water/ha
Oil Seed Crops			
	i. Canola	1 - 3 L 1 - 3 L	i. Apply 2 – 3 weeks after emergence ii. Apply 14 days later Apply in 200 L water/ha
	ii. Groundnuts	1 - 3 L 1 - 3 L	i. Apply 2 – 3 weeks after emergence ii. Apply 14 days later Apply in 200 L water/ha
	iii. Soy Beans	1 - 3 L 1 - 3 L	i. Apply 2 – 3 weeks after emergence ii. Apply 14 days later Apply in 200 L water/ha
	iv. Sunflower	1 - 3 L 1 - 3 L	i. Apply 2 – 3 weeks after emergence ii. Apply 14 days later Apply in 200 L water/ha
Ornamentals	Cut Flowers, Bulbs	1 - 3 L	i. Drench seedlings before transplant (see drench application below) ii. Apply 2 weeks after transplant iii. Repeat every 14 – 21 days Apply in 200 – 500 L water/ha
Grasses	Pastures and Lawns	3 - 5L 3 - 5L	i. Apply at the start of the growing season ii. Repeat application after 14 days Apply in 300 L water/ha
Bromeliaceae	Pineapples	1 - 3 L 1 - 3 L	i. Apply at the start of the growing season ii. Repeat application after 14 days Apply in 500 L water/ha
Pome fruit	Apples, Pears, Quinces	3 - 5L 3 - 5 L	i. Apply just before flowering ii. Apply at 100% petal fall iii. Apply 14 days later Apply in 1000 - 2000 L water/ha
Solanaceous Crops			
	i. Chilies, Paprika, Peppers,	1 - 2L	i. Drench seedlings before transplanting (see drench application below)
	Tomatoes	1 - 2.5L	ii. Apply 2 weeks after transplant iii. Repeat 14 days later, repeat 2 – 3times Apply in 200 L water
	ii. Tobacco	1 - 2 L 1 - 2,5 L	i. Drench seedlings before transplanting (see drench application below) ii. Apply 2 weeks after transplant iii. Repeat 14 days later, repeat 2 – 3times Apply in 200 L water
Stone Fruit	Apricots, Nectarines, Peaches, Plums Subtropical Fruit	3 - 5L 3 - 5 L	i. Apply just before flowering ii. Apply at 100% petal fall iii. Apply 14 days later Apply in 1000 - 2000 L water/ha
Sugar	Sugarcane	3 - 5 L 3 - 5 L	i. Apply at the start of the growing season ii. Repeat 2 – 3 times with 14 day intervals Apply in 300 L water/ha
Subtropical Fruit	i. Avocados	3 - 5L 3 - 5 L	i. Apply just before flowering ii. Apply at 100% petal fall iii. Apply 14 days later Apply in 1000 - 2000 L water/ha
	ii. Guavas	3 - 5L 3 - 5 L	i. Apply just before flowering ii. Apply at 100% petal fall iii. Apply 14 days later Apply in 1000 - 2000 L water/ha
	iii. Lychees, Mangoes	3 - 5L 3 - 5 L	i. Apply just before flowering ii. Apply at 100% petal fall iii. Apply 14 days later Apply in 1000 - 2000 L water/ha
	iv. Citrus	3 - 5L 3 - 5 L	i. Apply just before flowering ii. Apply at 100% petal fall iii. Apply 14 days later Apply in 1000 - 2000 L water/ha
Other	Other Crops	1 - 3 L 1 - 3 L	i. Apply at the start of the growing season ii. Repeat 2 – 3 times with 14 day intervals Apply in 300 L water/ha

APPLICATION METHODS

SOIL APPLICATION:		
All Crops	5 - 10 L	Apply through irrigation water or apply directly to the soil with suitable applicator. Use sufficient water to wash into root zone. Use the lower dosage on younger plants and higher dosage for mature plants. Repeat application every 14 – 21 days as necessary
DRENCH APPLICATION:		
Young Trees	2% Solution	Add 1 – 5 L solution as a soil drench or into the planting hole before planting.
Seedlings	2% Solution	Drench seedlings or seedling trays for 5-10 minutes in solution before transplanting.